

9.6 The Algebra of Functions

In this section, we will take a look at all our basic operations as they apply to functions.

Operations on Functions

If f and g are functions and x is an element of the domain of each function then,

$$(f + g)(x) = f(x) + g(x)$$

$$(fg)(x) = f(x)g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, g(x) \neq 0$$

This just means in order to perform a basic operation on two functions, you simply need to do that particular operation pointwise.

Example 1:

Let $f(x) = x^2 - x$ and $g(x) = 3x - 2$. Find the following.

a. $(f - g)(3)$

b. $(f + g)(2)$

c. $(fg)(-1)$

d. $(f + g)(x)$

e. $(fg)(x)$

f. $\left(\frac{f}{g}\right)(0)$

g. $\left(\frac{f}{g}\right)\left(\frac{2}{3}\right)$

Solution:

a. First we recall the rule given above $(f - g)(x) = f(x) - g(x)$.

There are two primary ways in which we can approach this problem. One way is to just replace the x in the rule with 3 and simplify as follows:

$$\begin{aligned}(f - g)(3) &= f(3) - g(3) \\ &= (3^2) - 3 - (3(3) - 2) \\ &= 9 - 3 - 9 + 2 \\ &= -1\end{aligned}$$

The alternate way to do this problem is to just find $(f - g)(x)$ in general and then substitute the value in the end.

$$\begin{aligned}(f - g)(x) &= f(x) - g(x) \\ &= x^2 - x - (3x - 2) \\ &= x^2 - 4x + 2\end{aligned}$$

$$(f - g)(3) = 3^2 - 4(3) + 2 = -1$$

We can see they both give us -1 . We prefer the latter method since once the expression for $(f - g)(x)$ has been found we can evaluate it at as many values as we would like.

b. Again we will start by finding $(f + g)(x)$ and then evaluate at 2.

$$\begin{aligned}
 (f+g)(x) &= f(x) + g(x) \\
 &= x^2 - x + (3x - 2) \\
 &= x^2 + 2x - 2 \\
 (f+g)(2) &= 2^2 + 2(2) - 2 = 6
 \end{aligned}$$

c.

$$\begin{aligned}
 (fg)(x) &= f(x)g(x) \\
 &= (x^2 - x)(3x - 2) \\
 &= 3x^3 - 5x^2 + 2x \\
 (fg)(-1) &= 3(-1)^3 - 5(-1)^2 + 2(-1) \\
 &= -10
 \end{aligned}$$

c. Since we have already found $(f+g)(x)$ for part b. we simply need to refer back.

$$(f+g)(x) = x^2 + 2x - 2$$

d. We have also already found $(fg)(x) = 3x^3 - 5x^2 + 2x$ in part c above.

e. We will again start by finding $\left(\frac{f}{g}\right)(x)$, then evaluate at 0.

$$\begin{aligned}
 \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \\
 &= \frac{x^2 - x}{3x - 2} \\
 \left(\frac{f}{g}\right)(0) &= \frac{0^2 - 0}{3(0) - 2} \\
 &= \frac{0}{-2} \\
 &= 0
 \end{aligned}$$

f. In part e. we found $\left(\frac{f}{g}\right)(x) = \frac{x^2 - x}{3x - 2}$. Evaluating at $\frac{2}{3}$ we get

$$\begin{aligned}
 \left(\frac{f}{g}\right)\left(\frac{2}{3}\right) &= \frac{\left(\frac{2}{3}\right)^2 - \left(\frac{2}{3}\right)}{3\left(\frac{2}{3}\right) - 2} \\
 &= \frac{\frac{4}{9} - \frac{2}{3}}{0}
 \end{aligned}$$

But we recall that we cannot have a zero in the denominator. Also, from the rules above, we know that $g(x) \neq 0$. Thus, we can see that $\frac{2}{3}$ is not in the domain of $\left(\frac{f}{g}\right)(x)$.

Thus, we can say $\left(\frac{f}{g}\right)\left(\frac{2}{3}\right)$ is not a real number.

We now turn our attention to another very important operation on functions, the composition.

Definition: Composition of two functions

Let f and g be two functions such that $g(x)$ is in the domain of f for all x in the domain of g . Then the composition of f and g , written $f \circ g$, is the function given by $(f \circ g)(x) = f(g(x))$.

It usually helps to visualize the composition as $f(g(x))$.

The composition idea is one of putting an entire function into another function.

Example 2:

Let $f(x) = 2x^2 - x + 1$, $g(x) = 2x - 3$ and $h(x) = \frac{1}{x+1}$. Find the following.

- a. $(f \circ g)(0)$ b. $(h \circ g)(2)$ c. $(h \circ f)(2)$ d. $(f \circ g)(x)$
 e. $(g \circ h)(x)$ f. $(f \circ f)(x)$

Solution:

- a. First, by the definition we know $(f \circ g)(x) = f(g(x))$. So we really want $f(g(0))$. Like the example above, we can do this by finding the formula for $(f \circ g)(x)$ first and then evaluate it at 0. So, if we insert $g(x) = 2x - 3$ into $f(g(x))$ and simplify we get

$$\begin{aligned}(f \circ g)(x) &= f(g(x)) \\ &= f(2x - 3) \\ &= 2(2x - 3)^2 - (2x - 3) + 1 \\ &= 2(4x^2 - 12x + 9) - 2x + 3 + 1 \\ &= 8x^2 - 24x + 18 - 2x + 4 \\ &= 8x^2 - 26x + 22\end{aligned}$$

Now we evaluate it for $x=0$

$$\begin{aligned}(f \circ g)(0) &= 8(0)^2 - 24(0) + 22 \\ &= 22\end{aligned}$$

- b. In a similar fashion we start by finding $(h \circ g)(x)$, then evaluate as follows

$$\begin{aligned}
 (h \circ g)(x) &= h(g(x)) \\
 &= h(2x-3) \\
 &= \frac{1}{(2x-3)+1} \\
 &= \frac{1}{2x-2} \\
 (h \circ g)(2) &= \frac{1}{2(2)-2} \\
 &= \frac{1}{2}
 \end{aligned}$$

c.

$$\begin{aligned}
 (h \circ f)(x) &= h(f(x)) \\
 &= h(2x^2 - x + 1) \\
 &= \frac{1}{(2x^2 - x + 1) + 1} \\
 &= \frac{1}{2x^2 - x + 2} \\
 (h \circ f)(2) &= \frac{1}{2(2)^2 - (2) + 2} \\
 &= \frac{1}{8}
 \end{aligned}$$

d. In part a. above we already found $(f \circ g)(x) = 8x^2 - 24x + 22$.

e.

$$\begin{aligned}
 (g \circ h)(x) &= g(h(x)) \\
 &= g\left(\frac{1}{x+1}\right) \\
 &= 2\left(\frac{1}{x+1}\right) - 3 \\
 &= \frac{2}{x+1} - 3
 \end{aligned}$$

e. Lastly, we want to take the composition of a function with itself. We do this in the same manner as all other compositions. We simply insert f into f and simplify.

$$\begin{aligned}
 (f \circ f)(x) &= f(f(x)) \\
 &= f(2x^2 - x + 1) \\
 &= 2(2x^2 - x + 1)^2 - (2x^2 - x + 1) + 1 \\
 &= 2(4x^4 - 4x^3 + 5x^2 - 2x + 1) - 2x^2 + x - 1 + 1 \\
 &= 8x^4 - 8x^3 + 10x^2 - 4x + 2 - 2x^2 + x \\
 &= 8x^4 - 8x^3 + 8x^2 - 3x + 2
 \end{aligned}$$

9.6 Exercises

Let $f(x) = 2x - 1$ and $g(x) = x^2 - 2x + 1$. Find the following.

1. $(f + g)(-1)$
2. $(f - g)(2)$
3. $(g - f)(x)$
4. $(fg)(-2)$
5. $\left(\frac{f}{g}\right)(0)$
6. $\left(\frac{g}{f}\right)(x)$
7. $(f - g)(a + h)$
8. $(f + g)(2 + h)$
9. $\left(\frac{f}{g}\right)(ab)$
10. $(fg)(x)$

Let $g(x) = \frac{1}{x+1}$ and $h(x) = \frac{x}{x+1}$. Find the following.

11. $(g + h)(1)$
12. $(g - h)(-2)$
13. $(gh)(x)$
14. $\left(\frac{g}{h}\right)(-2)$
15. $(h - g)(x)$
16. $(g + h)(x^2)$
17. $(h - g)(a - 1)$
18. $(gh)(0)$
19. $\left(\frac{h}{g}\right)(x)$
20. $(h - g)(a + b)$

Let $f(x) = \sqrt{x-1}$, $g(x) = x^2 + 1$ and $h(x) = \frac{2}{x}$. Find the following.

21. $(f + g)(x)$
22. $(g - g)(x)$
23. $(h - g)(x)$
24. $(g - h)(x)$
25. $(gh)(x)$
26. $(fh)(x)$
27. $(fg)(x)$
28. $\left(\frac{g}{f}\right)(x)$
29. $\left(\frac{g}{h}\right)(x)$
30. $\left(\frac{f}{h}\right)(x)$

Let $f(x) = 3x + 1$ and $g(x) = 2x^2 - x + 4$. Find the following.

31. $(f \circ g)(-1)$
32. $(f \circ g)(0)$
33. $(g \circ f)(2)$
34. $(g \circ f)(-3)$
35. $(f \circ g)(-4)$
36. $(g \circ f)(6)$
37. $(f \circ g)(x)$
38. $(g \circ f)(x)$
39. $(f \circ f)(x)$
40. $(g \circ g)(x)$

Let $g(x) = \frac{1}{x+1}$ and $h(x) = \frac{2}{x}$. Find the following.

- | | | | |
|----------------------|-----------------------|----------------------|----------------------|
| 41. $(g \circ h)(2)$ | 42. $(h \circ g)(-1)$ | 43. $(h \circ g)(0)$ | 44. $(g \circ h)(0)$ |
| 45. $(h \circ h)(2)$ | 46. $(g \circ g)(0)$ | 47. $(g \circ h)(x)$ | 48. $(h \circ g)(x)$ |
| 49. $(h \circ h)(x)$ | 50. $(g \circ g)(x)$ | | |

Let $f(x) = \sqrt{x+1}$, $g(x) = x^2 - 2$ and $h(x) = |2x - 1|$. Find the following.

- | | | | |
|--------------------------|--------------------------|--------------------------|---------------------------|
| 51. $(f \circ g)(2)$ | 52. $(g \circ f)(0)$ | 53. $(h \circ f)(3)$ | 54. $(g \circ h)(-1)$ |
| 55. $(f \circ h)(t)$ | 56. $(g \circ f)(c)$ | 57. $(g \circ f)(8 + a)$ | 58. $(f \circ g)(\Delta)$ |
| 59. $(g \circ g)(a - b)$ | 60. $(h \circ g)(a + b)$ | 61. $(g \circ f)(x + h)$ | 62. $(h \circ f)(x)$ |
| 63. $(g \circ f)(x)$ | 64. $(h \circ g)(x)$ | 65. $(g \circ h)(x)$ | 66. $(f \circ h)(x)$ |
| 67. $(f \circ g)(x)$ | 68. $(h \circ h)(x)$ | 69. $(g \circ g)(x)$ | 70. $(f \circ f)(x)$ |